

## Distribution of molluscan fauna in the Karangad estuarine mangroves, South East Coast of India

<sup>1</sup>Vellathi Venkatesan, <sup>1</sup>Chellappa Kalidas, <sup>2</sup>Pariyappanal U. Zacharia, and <sup>3</sup>Santhanam Rajagopal

<sup>1</sup>Regional Centre of Central Marine Fisheries Research Institute, Mandapam Camp, India.

<sup>2</sup>Central Marine Fisheries Research Institute, Tuticorin, India.

<sup>3</sup>Centre of Advanced Study in Marine Biology, Annamalai University, Parangipettai, India.

Corresponding author: V. Venkatesan, venkatcmfri@yahoo.co.in

**Abstract.** A survey has been made during February 2010 to know the pattern of molluscan fauna in the mangroves located along the Karangad estuary. During the present investigation in the research area, 25 species of molluscs were recorded. They belong to 14 genera, 10 families and 5 orders. In the study area, 13 species of gastropods namely, *Cerithidea fluviatilis*, *Terebralia palustris*, *Cerithium citrinum*, *C. scabridum*, *C. obeliscus*, *Littorina scabra*, *L. undulata*, *Planaxis sulcatus*, *Drupa margaritcola*, *D. heptagonalis*, *Thais rudolphi*, *T. bufo*, and *T. tissoti* and 12 species of bivalves – *Gafrarium tumidum*, *G. pectinatum*, *Crassostrea madrasensis*, *Macra cuneata*, *Tellina ala*, *T. bruguieri*, *Saccostrea cucullata*, *Modiolus metcalfei*, *M. tulipa*, *M. traillii*, *Meretrix meretrix* and *M. casta* - were recorded.

**Key Words:** mangrove, malacofauna, Karangad estuary, molluscs, India.

**Introduction.** The word "mangrove" is formed by combination of the Portuguese word "mangue" and the English word "grove". Mangroves, salt-tolerant plants, occur in most tropical and subtropical regions of the world. This group of plants is very important to the ecosystem diversity because they protect the coastline from destruction (maintain the ecosystem diversity) and provide many resources for utilization in the forestry, fisheries, food, agricultural and medicinal industries. The mangrove ecosystems are highly productive intertidal forests distributed along the tropical coast and they stabilize the coastal zone from erosion and act as a buffer zone between land and sea. Mangroves preserve water quality and reduce pollution by filtering suspended material and assimilating dissolved nutrients (Bandaranayake et al 2002).

Mangroves are predominantly intertidal habitats that occur along sheltered and shallow water coastlines. Mangrove-derived detritus is an important food source for decomposer food webs including many macro invertebrates (Fratini et al 2000; Cannicci et al 2008). The Pichavaram ecosystem of east coast of India is an estuarine-mangrove complex and supports a wide variety of biological species (Kathiresan 2000). Interacting with aquatic, inshore, upstream and terrestrial ecosystems, they support a diverse marine, freshwater and terrestrial flora and fauna (Macintosh & Ashton 2002) providing habitats for the distribution of diverse animals (Hogarth 2001). As such, mangrove roots become home to terrestrial as well as marine plants, algae, invertebrates and vertebrates. Molluscs are abundant in the littoral zones of sea. Molluscan group namely gastropods and bivalves form 98% of the total population. Macrobenthos may be operationally separated in three groups, i.e., epifauna, infauna and arboreal. Epifauna refers to those invertebrates that live on various substrates such as lower tree trunks and the sediment surface, but which do not burrow in it. Infauna refers to burrowing invertebrates which live within the sediment and arboreal forms refer to those live attached to stems, roots of the mangrove vegetation.

Many studies were carried out on the ecology and faunal distribution of mangrove swamps of Indo–West Pacific region (Macnae 1968), Malaysia (Brown 1971), west Indies (Coomans 1969), and in the province of Pavia, Northern Italy and North Western Australian mangrove swamps (Wells 1983). Mangrove faunal assemblages of molluscs relatively poorly known compared to other components of mangrove ecosystem. The investigation on the faunal assemblage of the molluscan fauna of the mangrove forests situated at the mouths of the estuary on the east coast of India is scanty. A survey has been made during February 2010 to know the pattern of molluscan fauna in the mangroves located in the karangad estuary.

## Materials and Methods

Karangad estuary is formed by the confluence of Kotai River with the sea on the Palk bay side at the southeast coast of India (Figure 1). The estuary lies between 9°38'42.1" N to 9°38'49.3" N and 78°57'54.2"E to 78°56'24.8" E. The width and depth of estuary ranges from 120m to 246m and from 1 to 2m respectively. This mangrove is bordered by two villages – Karangad and Pudukadu. The bottom of the estuary consists of clay, silt with admixture of sand and large amount of organic detritus.

The molluscan fauna of the extensive mangrove were collected by hand picking and the bivalve molluscs like mussels and oysters were collected by scrapping using knives from a known unit area of a quadrat (0.5m<sup>2</sup>). The infauna forms was collected by digging the substratum as described by (Alfred 1997).The arboreal forms were collected from the roots, stems and other parts of the mangrove trees vertically at every 25 cm height by hand picking (Sasekumar 1974).




Figure 1. Location of Karangad estuarine mangroves.

## Results and Discussions

A systematic list of molluscan fauna recorded in the karangad estuarine mangroves through out the study period is listed in Table 1.

**Mangroves as Habitats for Molluscs.** Molluscs can reach a high biomass in mangroves and they occupy very different levels of the ecosystem food web. Molluscs along with decapod crustaceans are the most well represented taxon of marine origin in mangrove forests (Plaziat 1984; Kathiresan & Bingham 2001). Molluscs dwell on the sediment surface or reside in burrows, others live on pneumatophores and lower tree trunks or prop-roots, burrow in decaying wood, or can even be found in the tree canopies (Sasekumar 1974; Ashton 1999). In mangroves, molluscs occupy all the levels of the food web, as predators, herbivores, detritivores and filter feeders. They are zoned both horizontally (i.e. along the sea-land axis) and vertically (i.e. at diverse heights from the ground) and include both mobile and sessile species. The presence of marine wood borers in the mangrove forest of Godavari estuary was reported by Ganapathi & Rao (1959). Studies on mangrove associated molluscan fauna of various Indian peninsular estuaries viz. Godavari and Krishna estuaries (Radhakrishnan & Janakiram 1975), Mahanadhi estuary (Subba Rao & Mookherjee 1975), Muriganga estuary, Sunderbans (Subba Rao et al 1983) have already been carried out. Murthy & Balaparameswara Rao (1977) carried out some works on the ecology of molluscs of the mangrove swamps in Machilipatnam. A survey on the investigation of malacofauna in the Pichavaram mangrove forest was done by Kasinathan & Shanmugam (1985). Hundred species of molluscs have been reported from mangroves areas of Andaman and Nicobar islands (Das & Roy 1989) and the mangrove fauna of Indian subcontinent (Dey 2006). A recent study on the malacofauna in Pondicherry mangroves was made by Khan et al 2008 and Kesavan et al 2009). Studies on molluscs overall consumption of mangrove litter in some Indo-Pacific mangrove forests using carbon and nitrogen stable isotope signatures and consequent contribution in nutrient dynamics was carried out by Bouillon et al (2002ab).

**Gastropods and Bivalves.** In mangrove ecosystem, gastropods contribute to entrap primary production within the food web by grazing fallen leaves and consuming mud (mainly formed by mangrove litter), bivalves are efficient filter feeders, able to capture suspended particles of various origins (Plaziat 1984; Kathiresan & Bingham 2001). Mud whelks consume significant amounts of fallen leaves in a Kenyan mangrove (five times the daily *R. mucronata* leaf production if fed ad libitum) was demonstrated by Fratini et al (2004). The impact of pollution in mangrove forests can be assessed based on the structure of the molluscan assemblages.

Mangrove roots and trunks represent zones of hard substrate colonised by fouling organisms e.g. Mussels and Oysters. Oysters, mussels and barnacles also foul mangrove roots and trunks (Pinto & Wignarajah 1980; Ross & Underwood 1997). Oyster and mussel form valuable fisheries in various parts of coastal India and are widely distributed in creeks, backwaters, bays and estuaries wherever hard substratum is found. The gastropod and bivalve molluscs are among the commonest epifaunal species that exist in the mangrove ecosystems. The epibenthic fauna includes primary and secondary consumers, and its impact on mangrove trees is considerable: barnacle assemblage can negatively affect root growth (Perry 1988) and heavy oyster cover can damage or break prop roots (Ellison & Farnsworth 1992, 2001). Gastropods and bivalves have a significant ecological role to play in the mangrove ecosystems and very little is known on their diversity of mangroves. Hence it is essential to document the biodiversity of ecosystems

In the present investigation, the recorded specimens were found to occur on mud banks, mud flats, mangrove forest, sandy muddy area swamps and hard substratum such as wooden poles and pillars. The oyster *Crassostrea madrasensis*, the clam *Meretrix casta* and the prosobranch gastropod *Cerethidia fluviatilis* and *Cerithium spp.* occurred by forming the beds in estuary. Oysters were found attached to wherever hard substratum is available such as wooden pole, wooden boat, mangrove stem, rock, brick, cement pillar or pole etc. Maximum number of species was collected from estuary banks and mud

flats. Most of the individuals were epifaunal and attached. *C. fluviatilis* were found to attach on to the mangrove stems and branches up to 1.0 meter height.

In the research area, the common species namely *T. palustris*; *C. scabridum*, *C. fluviatilis*, *C. obeliscus*, *Macra* spp. *L. scabra*, *T. ala*, *C. citrinum*, *T. bruguieri*, *M. casta* and *L. undulata* were found to occur in the muddy substratum. Certain species of Cerithium such as *C. citrinum*, *C. obeliscus*, *Modiolus* spp., *C. madrasensis*, *C. fluviatilis*, *Drupa* spp., *Thais* spp., and *Littorina* spp. were found to occur on the stem and root system of mangroves. *Gafrarium* spp. was found more near the mouth of estuary. *C. madrasensis* was found more in the middle of estuary by forming bed while *M. casta* was found more in the upper stream of the estuary. *Cerethidia* spp and *Cerithium* spp were forming beds through out the estuary and were predominant among all molluscan fauna.

Table 1 - A systematic list of molluscan fauna recorded in the Karangad estuarine mangroves

Taxonomical classification of molluscan fauna			
S. No.	Class : Gastropoda	S. No.	Class : Pelecypoda (bivalvia)
1.	Order : Mesogastropoda Family : Potamididae Genus : <i>Cerithidea</i> Species: <i>C. fluviatilis</i> (Potiez & Michaud)	1.	Order : Eulamellibranchiata Family : Veneridae Genus : <i>Meretrix</i> Species: <i>M. meretrix</i> (Linnaeus)
2.	Order : Mesogastropoda Family : Potamididae Genus : <i>Terebralia</i> Species: <i>T. palustris</i> (Linne)	2.	Order : Eulamellibranchiata Family : Veneridae Genus : <i>Meretrix</i> Species: <i>M. casta</i> (Chemnitz)
3.	Order : Mesogastropoda Family : Cerithiidae Genus : <i>Cerithium</i> Species: <i>C. citrinum</i> (Sowerby)	3.	Order : Mytiloidea Family : Mytilidae Genus : <i>Modiolus</i> Species: <i>M. metcalfei</i> (Hanley)
4.	Order : Mesogastropoda Family : Cerithiidae Genus : <i>Cerithium</i> Species: <i>C. scabridum</i> (Philippi)	4.	Order : Mytiloidea Family : Mytilidae Genus : <i>Modiolus</i> Species: <i>M. tulipa</i> (Lamarck)
5.	Order : Mesogastropoda Family : Cerithiidae Genus : <i>Cerithium</i> Species: <i>C. obeliscus</i> (Bruguere)	5.	Order : Mytiloidea Family : Mytilidae Genus : <i>Modiolus</i> Species: <i>M. traillii</i> (Reeve)
6.	Order : Mesogastropoda Family : Planaxidae Genus : <i>Planaxis</i> Species: <i>P. sulcatus</i> (Born)	6.	Order : Ostreoida Family : Ostreidae Genus : <i>Crassostrea</i> Species: <i>C. madrasensis</i> (Preston)
7.	Order : Neogastropoda Family : Muricidae Genus : <i>Drupa</i> Species: <i>D. margaritcola</i> (Broderrip)	7.	Order : Ostreoida Family : Ostreidae Genus : <i>Saccostrea</i> Species: <i>S. cucculata</i> (Born)
8.	Order : Neogastropoda Family : Muricidae Genus : <i>Drupa</i> Species: <i>D. heptagonalis</i> (Reeve)	8.	Order : Eulamellibranchiata Family : Veneridae Genus : <i>Gafrarium</i> Species: <i>G. tumidum</i> (Roding)

9.	Order : Neogastropoda Family: Muricidae Genus : <i>Thais</i> Species: <i>T. rudolphi</i> (Lamarck)	9.	Order : Eulamellibranchiata Family : Veneridae Genus : <i>Gafrarium</i> Species: <i>G .pectinatum</i> (Linne)
10.	Order : Neogastropoda Family : Muricidae Genus : <i>Thais</i> Species: <i>T. bufo</i> (Lamarck)	10.	Order : Eulamellibranchiata Family : Mactridae Genus : <i>Mactra</i> Species: <i>M. cuneata</i> (Chemnitz)
11.	Order : Neogastropoda Family : Muricidae Genus : <i>Thais</i> Species: <i>T. tissoti</i> (Petit)	11.	Order : Eulamellibranchiata Family : Tellinidae Genus : <i>Tellina</i> Species: <i>T. ala</i> (Hanley)
12.	Order : Mesogastropoda Family : Littorinidae Genus : <i>Littorina</i> Species: <i>L.(Littorinopsis) scabra</i> (Linnaeus)	12.	Order : Eulamellibranchiata Family : Tellinidae Genus : <i>Tellina</i> Species: <i>T. bruguieri</i> (Hanley)
13.	Order : Mesogastropoda Family : Littorinidae Genus : <i>Littorina</i> Species: <i>L. undulata</i> (Gray)		

**Conclusion.** During the present study, 25 species of molluscs (13 gastropods and 12 bivalves) were recorded. They belong to 14 genera, 10 families and 5 orders. The oyster *Crassostrea madrasensis*, the clam *Meretrix casta* and the prosobranch gastropod *Cerethidia fluviatilis* and *Cerithium spp.* occurred by forming the beds in estuary. Maximum number of species was observed from mud flats along the mangroves. Oysters were found attached to wherever hard substratum is available. *Cerethidia spp* and *Cerithium spp* were found predominant among the observed fauna forming bed throughout the estuary.

**Acknowledgements.** Authors are thankful to the director of Central Marine Fisheries Research Institute for providing the necessary facilities.

## References

- Alfred J. B., Varshney R. K., Ghosh A. K. (eds) 1997 An assessment manual for faunal biodiversity in South Asia. SACEP/NORAD Publication Series on Biodiversity in South Asia No **1**, 181pp.
- Ashton E. C., 1999 Biodiversity and community ecology of mangrove plants, molluscs and crustaceans in two mangrove forests in Peninsular Malaysia in relation to management practices. Ph.D. Thesis. University of York, UK.
- Bandaranayake W. M., 2002 Bioactivities, bioactive compounds and chemical constituents of mangrove plants. *Wetland Ecology & Management* **10**:421-452.
- Bouillon S., Koedam N., Raman A. V., Dehairs F., 2002b Primary producers sustaining macro-invertebrate communities in intertidal mangrove forests. *Oecologia* **130**:441-448.
- Bouillon S., Raman A. V., Dauby P., Dehairs F., 2002a Carbon and nitrogen stable isotope ratios of subtidal benthic invertebrates in an estuarine mangrove ecosystem (Andhra Pradesh, India). *Estuar Coast Shelf Sci* **54**:901-913.
- Brown D. A., 1971 Gonad development and spawning in *Anadara granosa* (L.)(Bivalvia: Arcidae). *Aquaculture* **30**:211-219.

- Cannicci S., Burrows D., Fratini S., Smith III T. J., Offenbergh J., Dahdouh-Guebas F., 2008 Faunistic impact on vegetation structure and ecosystem function in mangrove forests: A review. *Aquat Bot* **89**:186–200.
- Coomans H. E., 1969. Biological aspects of mangrove molluscs in the West Indies. *Malacologia* **9**:79-84.
- Das A. K., Roy M. K. D., 1989 A general account of the mangrove fauna of Andaman and Nicobar islands. Conservation Area Series, Zoological Survey of India.
- Dey A., 2006 Handbook on Mangrove Associate Molluscs of Sunderbans. Kolkata Z. S. I., (ed), 96p.
- Ellison A. M., Farnsworth E. J., 1992 The ecology of Belizean mangrove-root fouling communities: patterns of epibiont distribution and abundance, and effects on root growth. *Hydrobiologia* **247**:87–98.
- Ellison A. M., Farnsworth E. J., 2001 Mangrove communities. In: Bertness M. D., Gaines S., Hay M. E. (eds), *Marine Community Ecology*. Sinauer Press, Sunderland, pp.423-442.
- Fratini S., Cannicci S., Vannini M., 2000 Competition and interaction between *Neosarmatium meinerti* (Crustacea: Grapsidae) and *Terebralia palustris* (Mollusca: Gastropoda) in a Kenyan mangrove. *Mar Biol* **137**:309–316.
- Fratini S., Vigiani V., Vannini M., Cannicci S., 2004 *Terebralia palustris* (Gastropoda; Potamididae) in a Kenyan mangal: size structure, distribution and impact on consuming leaf litters. *Mar Biol* **114**:1173–1182.
- Ganapati P. N., Rao M. L. V., 1959 Incidence of marine borers in the mangrove of Godavari estuary. *Curr Sci* **28**(8):332.
- Hogarth P. J., 2001 Mangrove ecosystems. pp. 853-870. In: *Encyclopedia of Biodiversity*. Vol. **3**., Academic Press.
- Kasinathan R., Shanmugam A., 1985 Molluscan fauna of Pichavaram mangroves, Tamil Nadu. *Proc Nat Sym Biol Util Cons Mangroves, Kolhapur, Distribution, India*, 438-443.
- Kathiresan K., 2000 Flora and Fauna in Mangrove Ecosystems: A Manual for Identification. All India coordinated project on coastal and marine biodiversity, training and capacity building on coastal biodiversity (east coast), Ministry of Environment and Forests, CAS in Marine Biology, Parangipettai, India.
- Kathiresan K., Bingham B. L., 2001 Biology of mangroves and mangroves ecosystems. *Adv Mar Biol* **40**:84–251.
- Kesavan K., Baabu A., Ravi V., Rajagopal S., 2009 A checklist of malacofauna from Pondicherry mangroves. *AES Bioflux* **1**(1):31-36.
- Khan A. B., Saravanan K. R., Ilangovan K., 2008 Floristic and macro faunal diversity of Pondicherry mangroves, South India. *Tropical ecology* **49**(1):91-94.
- Macintosh D. J., Ashton E. C., 2002 A review of mangrove biodiversity conservation and management. Final Report 10/06/2002. Centre for Tropical Ecosystems Research, University of Aarhus, Denmark.
- Macnae W., 1968 A general account of the fauna and flora of mangrove swamps and forests in the Indo-West Pacific region. *Adv Mar Biol* **6**:73-270.
- Murty A. S., Balaparameswara Rao M., 1977 Studies on the ecology of mollusks in a South Indian Mangrove swamp. *J Moll Stud* **43**:223–229.
- Perry D. M., 1988 Effects of associated fauna on growth and productivity in the red mangrove. *Ecology* **69**:1064–1075.
- Pinto L., Wignarajah S., 1980 Some ecological aspects of the edible oyster *Crassostrea cucullata* (Born) occurring in association with mangroves in Negombo lagoon, Sri Lanka. *Hydrobiologia* **69**:11–19.
- Plaziat J. C., 1984 Mollusk distribution in the mangal. In: Por D., Dor I. (eds), *Hydrobiology of the Mangal: the Ecosystem of the Mangrove Forests*. Junk, Boston, pp. 111–143.
- Radhakrishna Y., Janakiram R., 1975 The mangrove molluscs of Gadavari and Krishna estuary. In: *Recent researches in estuarine biology*, Natarajan R. (ed), Hisdustan publishing corporation (L), Delhi, India, pp. 177-184.

- Ross P. M., Underwood A. J., 1997 The distribution and abundance of barnacles in a mangrove forest. *Aust J Ecol* **22**:37-47.
- Sasekumar A., 1974 Distribution of the macro fauna on a Malayan mangrove shore. *J Anim Ecol* **43**:51-69.
- Subba Rao N. V., Mookherjee H. P., 1975 On the collection of molluscs from Mahanadi estuary, Orissa. In: *Researches in estuarine biology*, Natarajan R. (ed), 165-176. Hindustan publishing corporation (L), Delhi, India.
- Subba Rao N. V., Dey A., Baruna S., 1983 Studies on the malacofauna of Muriganga estuary, Sunderbans, West Bengal. *Bull Zool Surv India* **5**(1):47-56.
- Wells F. E., 1983 An analysis of marine invertebrate distribution in a mangrove swamp in northwestern Australia. *Bulletin of Marine Science* **33**:736-744.

Received: 10 August 2010. Accepted: 11 September 2010. Published online: 13 September 2010.

Authors:

Vellathi Venkatesan, Molluscan Fishery Division, Regional Centre of Central Marine Fisheries Research Institute, Mandapam Camp 623520, Tamil Nadu, India, e-mail: venkatcmfri@yahoo.co.in

Chellappa Kalidas, Mariculture Division, Regional Centre of Central Marine Fisheries Research Institute, Mandapam Camp 623520, Tamil Nadu, India, e-mail: dascift@gmail.com

Pariyappanal Ulahannan Zacharia, Demersal Fishery Division<sup>2</sup>Central Marine Fisheries Research Institute, Tuticorin 628001, India. e-mail: Zachariapu@yahoo.com

Santhanam Rajagopal, Centre of Advanced Study in Marine Biology, Annamalai University, Parangipettai 608502, India. e-mail: rajgopi52@yahoo.com

How to cite this article:

Venkatesan V., Kalidas C., Zacharia P. U., Rajagopal S., 2010 Distribution of molluscan fauna in the Karangad estuarine mangroves, South East Coast of India. *AES Bioflux* **2**(2):113-119.

